

## Suprep 2 Day Preparation

Procedure is scheduled with: \_\_\_\_\_

Date: \_\_\_\_\_

Arrival time: \_\_\_\_\_

Procedure time: \_\_\_\_\_

### Procedure Location

Out Patient Endoscopy Locations			
<b>Clayton</b> 900 S Lombard Street. Suite# 104  Phone: 919-341-3547	<b>Raleigh</b> 2601 Lake Drive Suite# 201  Phone: 919-783-4888	<b>Wake Forest</b> 10540 Ligon Mill Road Suite# 109  Phone: 919-439-3393	<b>Wilson</b> 2402 Camden St SW Suite #300  Phone: 252-237-4100
<b>Out Patient at Wake Med Hospital</b> Cary                      North                      New Bern		<b>Out Patient at Rex Hospital</b>	

**You will be sedated for this procedure and you must have:**

- **Adult driver, 18 years or older to bring you to your procedure**
- **Adult driver must stay at the facility for the duration of the procedure**
- **Adult driver must drive you home**

**If you take Phentermine it MUST be stopped 10 days prior to procedure**

**If you take coumadin, warfarin, pradaxa, or any other blood thinner please call our office for specific instructions**

The facility has the right to delay or cancel your procedure in the event you do not have an adult with you to drive you home.

If you are unable to keep your appointment, please call our office as soon as possible to reschedule. We do reserve the right to charge for missed appointments or appointments cancelled less than 72 hours in advance.

### Diabetic Patients

If taking diabetic medication:

- Day before procedure:  
AM --Take your normal insulin and/or oral medications.  
PM --Take ½ of PM insulin dose, NO oral diabetic medication.
- Day of procedure:  
Do not take any diabetic medications until after the procedure.

### Asthma Patients

If using an in haler

- Bring inhaler with you on day of procedure

### CPAP or VPAP

If you use either of these machine, you are required to bring it with you.

## Suprep 2 Day Preparation for Colonoscopy

1). **Follow instructions outlined below.** Please do not follow instructions found in the Suprep kit received from the pharmacy.

**Bowel prep will include the following items sent to your pharmacy:**

- 1-Suprep Kit,
- 2-Reglan tablets (generic name is Metoclopramide)
- 2-Dulcolax tablets (generic name is Bisacodyl) this can be purchased over the counter, no prescription required)

2). **5 days prior**

- Stop any iron supplements you are taking.
- Avoid sesame seeds and nuts

3). **Day before procedure:**

- Drink clear liquids **only for all snacks and meals.**  
Clear liquid diet: coffee, tea, soft drinks, Jell-O, clear broth, popsicles, clear fruit juices, Gatorade, water, Snapple, powdered juices. Avoid red or purple liquids
- Force fluids all day prior to and after beginning the bowel prep to keep you hydrated
- No solid foods or milk products

**2:00 pm:** Take the 2 Dulcolax tablets.

**5:30pm:** Take the Reglan tablet (this is optional). The prep may cause nausea. Taking the Reglan in advance of the prep should help reduce the nausea.

**6:00 pm:** Pour ONE of the 6 ounce bottles of Suprep liquid in the container provided.  
Add cool drinking water up to the 16 ounce line of the container.  
Drink **all** liquid in the container.  
Drink 2 additional 16 ounces of water over the next hour

~Do not plan to go anywhere after you have started drinking the prep.

The prep will cause diarrhea to cleanse your bowel for the procedure.

~Using baby wipes rather than toilet tissue to protect your rectum from irritation.

**DO NOT eat or drink anything after midnight other than the 2<sup>nd</sup> prep dose as instructed below**

**4) Morning of procedure -- 2<sup>nd</sup> Dose**

**Start so you are finished 4 hours prior to arrival time.**

- 30 minutes before starting 2nd dose take 1-Reglan tablet.  
Pour remaining 6 ounce bottles of Suprep liquid in the container provided.  
Add cool drinking water up to the 16 ounce line of the container.  
Drink **all** liquid in the container.  
Drink 2 additional 16 ounces of water over the next hour
- Do not drink anything after you have completed the prep
- Do not eat anything until after the procedure